

Chlorothalonil Application and Use Chart

| Crop | Current Label | PMRA Final Decision | | | | |
|--|---------------|--|-------------------------------|---|----------------------|------------------------------|
| | # apps | # apps | Application rate (kg a.i./ha) | Max area/day if mixing and loading using non-closed and ground-boom application method (max. 340 kg a.i./day) | | Re-treatment Interval (days) |
| | | | | Hectares | Acres | |
| Asparagus | 3 | 3 | 1.7 (SN) 1.2 (DF) | 200 (SN) 283 (DF) | 494 (SN) 699 (DF) | 14 |
| Blueberries (HB) | 3 | 2 | 3.6 | 94 | 232 | 7 |
| Blueberries (LB) | 2 | 2 | 3.6 (SN) 2.5 (DF) | 94 (SN) 136 (DF) | 232 (SN) 336 (DF) | 42 |
| Cabbage | 5 | 2 | 2.4 | 141 | 348 | 7 |
| Carrot | 7 | 7 | 1.6 | 212 | 524 | 7 |
| Celery, field | 9 | 2 | 2 | 170 | 420 | 3 |
| Celery seedbeds (greenhouse) | 9 | 1 | 1.4 | 242 | 598 | N/A |
| Cherries (sweet and sour) | 4 | 3 (2 spring + 1 post-harvest) | 4.5 | 75 | 185 | 10 |
| Chickpeas | 3 | 2 (1 st at 2.0 kg a.i./ha and 2 nd at 1.5 kg a.i./ha) | 2.0 – 1 st | 170 | 420 | 10 |
| | | | 1.5 – 2 nd | 226 | 558 | |
| Cole crops (broccoli, Brussels sprouts, cauliflower) | 5 | 1 | 2.4 | 141 | 348 | N/A |
| Corn, sweet | 2 | 2 | 1.6 | 212 | 524 | 10 |
| Cranberries | 3 | 1 | 5.8 | 58 | 143 | N/A |
| Cucurbit vegetables (cantaloupe, muskmelon, honeydew, squash, pumpkin, watermelon, cucumber) | 7 | 2 | 2.4 | 141 | 348 | 7 |
| Ginseng | 6 | 3 (2 + 1 fall) | 2.4 | 141 | 348 | 7 |
| Hazelnut | 3 | 3 | 3.4 | 100 | 247 | 20 |
| Lentils | 2 | 2 | 2 | 170 | 420 | 10 |
| Mushrooms | 2 | 1 | 12.7 | 26 | 64 | N/A |
| Onion (dry bulb) | 3 | 2 | 2.4 | 141 | 348 | 7 |
| Onion (green bunching) | 5 | 2 | 2.4 | 141 | 348 | 7 |
| Parsnip | 7 | 7 | 1.4 | 242 | 598 | 7 |
| Pea, dry | 3 | 2 | 1.5 | 226 | 558 | 10 |

Continued on next page →

| Crop | Current Label | PMRA Final Decision | | | | |
|-----------------------------|---------------|---|-------------------------------|--|-------|--------------------------------|
| | # apps | # apps | Application rate (kg a.i./ha) | Max area/day if mixing and loading using non-closed and groundboom application method (max. 340 kg a.i./day) | | Re-treatment Interval (days) |
| | | | | Hectares | Acres | |
| Peach, nectarine | 4 | 3 (2 spring + 1 dormant) | 4.5 | 75 | 185 | 10 |
| Potato (seed) | 12 | 3 | 1.2 | - | - | 7 |
| Potato (table) | 12 | 3 | 1.2 | - | - | 7 |
| Strawberries | 3 | 3 (2 spring + 1 post-harvest) | 1.8 | 188 | 464 | 10 |
| Tomato (not for processing) | 9 | 2 total | 2.4 or | 141 | 348 | 14 when used at 2.4 kg a.i./ha |
| | | | 1.2 | 283 | 699 | 8 when used at 1.2 kg a.i./ha |
| Tomato (for processing) | 9 | 2 (2.4 kg a.i./ha) + 7 (1.2 kg a.i./ha) | 2.4 | 141 | 348 | 14 when used at 2.4 kg a.i./ha |
| | | | 1.2 | 283 | 699 | 8 when used at 1.2 kg a.i./ha |
| Wheat | 3 | 2 | 1.3 | 261 | 645 | 10 |

SN: Solution

DF: Dry formulations (Dry Flowable or Water Dispersible Granules) and all they have to be **re-formulated in WSP** (Water Soluble Packaging)